

The Actin Binding Domain of β I-Spectrin Regulates the Morphological and Functional Dynamics of Dendritic Spines

Michael W. Nestor^{1,2,3,ma}, Xiang Cai¹, Michele R. Stone^{1,mb}, Robert J. Bloch^{1,2,3}, Scott M. Thompson^{1,2,3,*}

1 Department of Physiology, University of Maryland School of Medicine, Baltimore, Maryland, United States of America, **2** Program in Neuroscience, University of Maryland School of Medicine, Baltimore, Maryland, United States of America, **3** Training Program in Integrative Membrane Biology, University of Maryland School of Medicine, Baltimore, Maryland, United States of America

Abstract

Actin microfilaments regulate the size, shape and mobility of dendritic spines and are in turn regulated by actin binding proteins and small GTPases. The β I isoform of spectrin, a protein that links the actin cytoskeleton to membrane proteins, is present in spines. To understand its function, we expressed its actin-binding domain (ABD) in CA1 pyramidal neurons in hippocampal slice cultures. The ABD of β I-spectrin bundled actin in principal dendrites and was concentrated in dendritic spines, where it significantly increased the size of the spine head. These effects were not observed after expression of homologous ABDs of utrophin, dystrophin, and α -actinin. Treatment of slice cultures with latrunculin-B significantly decreased spine head size and decreased actin-GFP fluorescence in cells expressing the ABD of α -actinin, but not the ABD of β I-spectrin, suggesting that its presence inhibits actin depolymerization. We also observed an increase in the area of GFP-tagged PSD-95 in the spine head and an increase in the amplitude of mEPSCs at spines expressing the ABD of β I-spectrin. The effects of the β I-spectrin ABD on spine size and mEPSC amplitude were mimicked by expressing wild-type Rac3, a small GTPase that co-immunoprecipitates specifically with β I-spectrin in extracts of cultured cortical neurons. Spine size was normal in cells co-expressing a dominant negative Rac3 construct with the β I-spectrin ABD. We suggest that β I-spectrin is a synaptic protein that can modulate both the morphological and functional dynamics of dendritic spines, perhaps via interaction with actin and Rac3.

Citation: Nestor MW, Cai X, Stone MR, Bloch RJ, Thompson SM (2011) The Actin Binding Domain of β I-Spectrin Regulates the Morphological and Functional Dynamics of Dendritic Spines. PLoS ONE 6(1): e16197. doi:10.1371/journal.pone.0016197

Editor: Lin Mei, Medical College of Georgia, United States of America

Received: August 24, 2010; **Accepted:** December 7, 2010; **Published:** January 31, 2011

Copyright: © 2011 Nestor et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Supported by stipends to MWN from T32 GM08181, to MRS from T32 NS07375, and from grants from the Muscular Dystrophy Association (RJB) and the National Institutes of Health (R01 HL75106 to RJB and R01 MH65488 to SMT). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: sthom003@umaryland.edu

^{ma} Current address: National Institute of Child Health and Development, National Institutes of Health, Bethesda, Maryland, United States of America

^{mb} Current address: Aeras Global TB Vaccine Foundation, Rockville, Maryland, United States of America

Introduction

Most excitatory synapses are positioned on dendritic spines, which display a range of sizes and shapes. Changes in dendritic spine size accompany synapse maturation [1] and are correlated with synaptic strength [2–4]. Spine morphology is altered by changes in synaptic activity [5,6] and can accompany changes in synaptic strength in some, but not all, forms of long-term potentiation [7–9]. Conversely, abnormal dendritic spines are observed in many forms of mental dysfunction. The signaling mechanisms that regulate spine structure and couple changes in synaptic structure and function are now being actively explored.

Actin, both G-actin monomers and filamentous, polymerized F-actin, is highly concentrated in dendritic spines [10] and regulates spine morphology. Dynamic changes in actin polymerization underlie morphological plasticity of spines [11,12] and can affect synaptic function [10,13–17]. Actin also serves as the scaffold to which transmembrane proteins like AMPA and NMDA receptors, as well as intracellular signaling molecules, are anchored [18–20]. Treatment with actin-depolymerizing agents reduces the number

of functional AMPARs at dendritic spines [21,22]. Thus, actin and its binding proteins, as well as the signaling molecules that regulate their interaction, are essential for synapse structure and function.

Spectrin is a large protein that links the actin cytoskeleton to the cytoplasmic surface of plasma membranes [23,24] and regulates the dynamic state of the actin cytoskeleton [25]. Spectrin also anchors transmembrane proteins, such as NMDARs [26], to the actin cytoskeleton, generating localized microdomains that facilitate rapid signaling cascades. Spectrin is a heterodimer, consisting of α - and β -chains [27,28]. β I-spectrin is enriched in postsynaptic spine heads in the absence of α -subunits [29]. Proteins in the spectrin superfamily are composed of a series of triple helical repeats that serve as binding sites for other cytoskeletal proteins. Spectrin also harbors a central SH3 domain and binding sites for Ca^{2+} -activated proteins, suggesting that it may play a role in signaling cascades. The actin-binding domain (ABD) is limited to the N-terminal region of β I-spectrin and is composed of two calponin homology domains.

As a first step in characterizing the role of spectrin in dendritic spines, we transfected CA1 pyramidal neurons in hippocampal

slice cultures with the ABD of spectrin or closely related proteins, following the approach previously used to study α -actinin [30]. We suggest that β I-spectrin may play both a structural and functional role in dendritic spines through interaction with the actin cytoskeleton and Rac3.

Methods

Ethical approval

All experimental protocols that involve tissue collected from animals in this study have been approved by The University of Maryland School of Medicine Institutional Animal Care and Use Committee (Protocol #0609015).

Neuronal cultures

Organotypic hippocampal slice cultures were prepared with the roller tube method [31]. Briefly, hippocampi were dissected from 5–7 day-old rat pups and cut into 400 μ m-thick slices with a chopper, under sterile conditions. Slices were attached to cleaned, poly-lysine-coated coverslips in a mixture of lyophilized chicken plasma (Cocalico, Reamstown, PA) and concentrated fibrin solution (Tisseel, Baxter Health Care) that was then clotted with thrombin. Coverslips containing slices were placed in culture tubes on a slowly rotating roller drum in an incubator. After 4 days *in vitro*, cultures were treated with antimetabolites overnight to reduce glial proliferation. Cultures remained in the incubator for 12–16 days before transfection. Cortical neuronal cultures were prepared and cultured from E18 rat neocortex, as described [12].

ABD and Rac constructs

Plasmids were created using PCR to amplify the cDNAs of the actin-binding domains (ABDs) of dystrophin, utrophin, actinin-2, and β I-spectrin [detailed methods: 32]. The first 942 base pairs of the coding sequence of the ABD of β I-spectrin (NM_000347) was amplified from plasmid B259 [32]. With primers A, 5' CGTAGAAT TCCTATG-ACATCGGCCACAG 3' (sense), and B, 5' TCCCC-GCGGCTAAGGTGAG CAGGTCCGA 3' (antisense). cDNA was cloned into the DsRed2-c1 plasmid vector (BD Biosciences Clontech) via a EcoRI sense site and a SacII antisense site. For dystrophin, a plasmid (pDys246) [33] was used with primers A, 5' GCGAATTC-TATGTTGTTGTGGGAAGAAGTA 3' (sense), and B, 5' GCAGGTACCCTATTC AATGCTC ACTTGTGTTG 3' (antisense), to amplify and clone the first 738 bases of the dystrophin gene (accession no. M18533), containing its ABD. The dystrophin ABD was cloned into the DsRed2-c1 vector via an EcoRI site on the sense primer and a KpnI site on the antisense primer. An utrophin plasmid (pet23Utr261) [34] served as a template to amplify the ABD. The primers A, 5' TATAAGCTTC-TATGGCCAAGTATGGGAC 3' (sense) and B, 5' TGTGGATCCCCTAATCTATCGTGACTT-GCTG 3' (antisense) were used to amplify the first 783 bases of the utrophin gene (accession no. Y12229), which was then cloned into the DsRed2-c1 vector via a HindIII sense site and a BamHI antisense site. The ABD of α -actinin-2 (CAA32078.1) was cloned into the DsRed2-c1 vector similarly, with the primers. A, 5' GTA-GAATTCATG AACACAGCATGAACCAG 3' (sense) and B, 5' TGTGGATCCCCTAATTCAGCAATTTCCATG 3' (antisense) were used to amplify the first 671 bases of the α -actinin-2 gene, via an EcoRI sense site and a BamHI antisense site. GFP-actin was from Clontech (Mountain View, CA). cDNAs encoding wild type and dominant negative Rac 1 and 3 were myc- and HA-tagged constructs from the Guthrie cDNA Resource Center (Rollo, MO). Plasmids were introduced into bacteria and amplified using standard protocols. Amplified cDNA was isolated and purified using a HiSpeed Plasmid Maxi Kit (Qiagen; Valencia, CA) following the

manufacturer's instructions and prepared for the biolistic transfection of organotypic hippocampal slice cultures.

Biolistic transfection

Cultures were transfected biolistically with a gene gun (Helios; Bio-Rad, Hercules, CA) as described [35]. Gold pellets (1.0 μ m diameter) were coated with spermidine and then placed in a solution of 25 μ g/ μ l DNA, which was precipitated onto the particles by the addition of CaCl₂ to a final concentration of 500 mM. The coated pellets were attached to the walls of the cartridges with polyvinylpyrrolidone, as described. Prior to transfection, each slice was placed in a small volume of solution containing 40 μ M amino-5-phosphonovaleric acid (AP5), 40 μ M 6,7-dinitroquinoxaline-2,3-dione (DNQX), and 10 mM Mg²⁺ to block synaptic transmission and reduce excitotoxicity. Cultures were transfected by shooting the gold particles out of the gene gun through a nylon mesh (90 μ m² pore size) at a distance of 2 cm from the culture surface at a pressure of 200 psi. After transfection, slices were returned to roller tubes and placed in the incubator for 24–48 hrs before being used for electrophysiology or imaging.

Electrophysiology

Cultures were placed in a recording chamber and perfused with extracellular saline containing (in mM): 145 NaCl, 10 NaHCO₃, 2 CaCl₂, 2 MgCl₂, and Phenol Red (10 mg/l). Transfected neurons were identified by fluorescence microscopy before recordings were performed. Patch pipettes were filled with the following (in mM): 130 gluconic acid, 10 KCl, 1 MgCl₂, 2 ATP, 10 HEPES, and 0.1 EGTA, titrated to pH 7.4 with KOH. To ensure that transfected neurons were those being recorded from, the pipette solution also contained Alexa 488 (100 μ M). After breakthrough to whole-cell configuration, transfected neurons contained both green Alexa 488 and red DsRed fluorescence, whereas untransfected cells, recorded within the same slice culture, displayed only green fluorescence. Miniature excitatory postsynaptic currents (mEPSCs) were recorded at a holding potential of -70 mV in the presence of tetrodotoxin (1 μ M) and bicuculline (40 μ M) at room temperature. Recording pipette resistances were between 6.5 and 8 M Ω . Recordings in which the access resistance exceeded 30 M Ω were discarded. Spontaneous mEPSCs were amplified using an Axopatch 200B amplifier and Clampex 9 software (Molecular Devices, Sunnyvale, CA), low-pass filtered at 2 kHz, and digitized at 10 kHz. mEPSCs were identified and characterized using a template-matching algorithm created from the average of 50 events. To be included for analysis, mEPSCs had to have a uniphasic rise time of <6 ms, a decay time constant of <25 ms and a peak amplitude >4 pA. Decay time constants were calculated by fitting each event with a single exponential.

Microscopy and data analysis

In experiments involving static imaging, spine head width and spine length were measured from confocal micrographs of all morphological spine categories. Confocal images were taken with an upright Zeiss LSM-510 Meta Confocal Microscope System using a water immersion objective (Zeiss; 60X/NA 1.2, C-Apochromat, effective pixel size = 0.017 μ m²). Fluorescence excitation was achieved using a 25 mW Argon ion laser (Spectra-physics, Mountain View, CA) and illuminating the specimen using the 488 nm and 514 nm lines.

3D image stacks were collected from living cells in unfixed cultures, background fluorescence was subtracted, and images were collapsed into maximal fluorescence 2D projections. Spines were then traced manually and the spine head width or the spine length was computed using the SimplePCI software with the

experimenter blinded to the condition. Spine head width was defined as the maximum dimension of the spine head, whereas spine length was defined as the distance from the base of the spine neck to the tip of the spine head. Spine density was calculated by manually counting the number of spines expressed over a 30 μm segment of dendrite. For all measurements of spine density or dimensions, at least 2 well isolated apical dendrites were used per neuron. In experiments involving dynamic imaging, these same analyses were performed on images taken from living cells in unfixed cultures collected using an epifluorescence wide-field microscope (Nikon; 60X/0.8 NA, water immersion objective) and a CCD camera (Hamamatsu Orca ER; effective pixel size = 0.012 μm^2). Photodamage was minimized by adding Trolox (0.2 mM) to the saline.

For post-hoc analysis of time-lapse image sequences, spine heads were traced manually and the length and width were computed. This method underestimates the morphological dynamics of spines because the measurements are only taken in x and y planes [36]. Images were captured every second for 3 min and binned into 10 s intervals. Morphological dynamics were measured by calculating the coefficient of variation of the spine head diameter within each 10 s interval. Out-of-focus images were removed from the analyses. Fluorescence intensity was calculated after subtracting the average background intensity of two regions of interest surrounding the dendritic spine.

To measure PSD-95 area, slice cultures were co-transfected biolistically with GFP fused to PSD-95 and DsRed fused to the β I-spectrin ABD. PSD-95 - GFP was a gift from Dr. T. A. Blanpied, and was constructed by subcloning PSD-95 at HindIII-EcoRI into N1 versions of the fusion protein expression constructs. After 24–48 h, slices were placed in a standard recording chamber filled with control saline and wide-field images were taken, as above. After background subtraction, each cluster of PSD-95-GFP was traced manually and an area was calculated using SimplePCI software.

Analyses of dendritic spine dimensions and PSD-95-GFP area were performed by comparing the cumulative probability distributions with the Kolmogorov-Smirnov test. Otherwise, unpaired t-tests were used to establish significance. When comparing more than two groups, ANOVA was used with a Tukey's or Scheffe's post-hoc analysis to establish significance unless otherwise noted. All values are presented as mean values \pm standard error of the mean.

Immunoprecipitation

Cultures of cortical neurons were washed with buffered saline with complete protease inhibitors (Roche Applied Science, Indianapolis, IN), placed in 0.32 M sucrose in the same solution, and scraped. After brief centrifugation, the cellular pellet was mixed with an equal volume of two-fold concentrated RIPA buffer to obtain the following final concentrations of detergents: 1% NP-40, 0.5% deoxycholate, 0.1% SDS. Extracts were triturated in the cold, briefly, and stored on ice. Aliquots (70 μl) of Dynabeads (Invitrogen, Carlsbad, CA), pre-coated with sheep anti-rabbit IgG, in buffered saline were bound with 6 μg affinity-purified rabbit antibodies to β I-spectrin or with purified non-immune rabbit IgG and washed. The bead pellet was mixed with 1 mg of protein of the cellular extract and mixed gently overnight. The immunoprecipitates were collected, washed briefly in buffered saline containing 0.5% Tween, solubilized with SDS-PAGE sample buffer, and subjected to SDS-PAGE, followed by electro-transfer to nitrocellulose. Blots were incubated with antibodies to β I-spectrin, Rac1, or Rac3. Bound antibody was detected with peroxidase-conjugated secondary antibodies (Roche Diagnostics,

Indianapolis, IN; Amersham Biosciences, England) and the SuperSignal West Femto Kit (Thermo Scientific, Waltham, MA). Samples of the cell extract were run in parallel, to mark the positions of the proteins of interest. Affinity-purified rabbit antibodies to β I-spectrin have been described [37]. Antibodies to Rac1 were from Santa Cruz Biotechnology (Santa Cruz, CA). Antibodies to Rac3 were the generous gift of Dr. I. De Curtis (San Raffaele Scientific Inst., Milan, Italy).

Results

The ABD of β I-spectrin induces actin bundling in CA1 pyramidal neurons

To investigate the interaction between spectrin family proteins and the actin cytoskeleton, we created constructs of the highly homologous ABDs of β I-spectrin, α -actinin-2, utrophin, dystrophin, and filamin (supporting information Fig. S1), all fused to the fluorophore DsRed, and introduced them biolistically into CA1 pyramidal cells in hippocampal slice cultures. We co-transfected with a GFP plasmid in most experiments to fill the cells and allow visualization of dendrites and dendritic spines (Fig. 1A), and quantified spine shape and size from images of GFP distribution.

The DsRed-tagged β I-spectrin ABD labeled filamentous structures in the cell body and apical and basal dendrites ($n = 12$)(Fig. 1B). We observed a distribution of the ABD of β I-spectrin that was similar to what has been reported for endogenous spectrin in hippocampal pyramidal neurons [29]. Co-transfection of GFP-tagged actin revealed that these structures were bundles of filamentous actin, which were not seen with expression of actin-GFP alone, suggesting that the ABD of β I-spectrin associates with actin filaments, promoting their formation and inducing bundling. The ABD of β I-spectrin, either alone or together with actin-GFP, also labeled spine heads in apical and basal dendrites (Fig. 1E).

Expression of the ABD of other members of spectrin superfamily tagged with DsRed gave different results. Expression of the ABD of utrophin resulted in actin bundling in the neuronal cell bodies, as well as apical and basal dendrites ($n = 12$)(Fig. 1C), but minimal labeling of dendritic spines (Fig. 1F). Expression of the ABDs of actinin-2 ($n = 6$) and filamin ($n = 7$), in contrast, produced diffuse labeling of the soma and principal dendrites (Fig. 1D), without actin bundling, as well as prominent labeling of dendritic spines (Fig. 1F). The ABD of dystrophin ($n = 12$) produced no actin bundling in the cell bodies or dendrites and did not label spines (Fig. 1F). Co-transfection of GFP with the utrophin and dystrophin constructs revealed that spines were present, but unlabeled by their ABDs. We observed that the expression level for all of the ABD constructs were similar based on the distribution and fluorescence intensity of each ABD. These results demonstrate that the ABDs of spectrin superfamily members caused distinct effects on actin filaments and were targeted differentially to dendritic spines. Although these ABDs are highly homologous, their effects are therefore likely to be mediated by more than mere binding to actin.

The ABD of β I-spectrin enlarges dendritic spine heads

Endogenous β I-spectrin is expressed at high levels in the heads of postsynaptic spine [29]. Analysis of confocal images of GFP-filled dendritic spines revealed that the heads of dendritic spines co-expressing the ABD of β I-spectrin were more than twice as large as spines expressing GFP alone ($n = 500$ spines/5 neurons/group; $p < 0.05$, K-S test)(Fig. 2). Spine length was not significantly different between the two groups, however ($n = 500$ spines/5 neurons/group, $p > 0.05$, K-S test)(Fig. 2B). Neurons co-expressing the ABD of β I-spectrin and GFP also displayed a small ($\sim 20\%$),




Figure 1. Actin binding domains from various spectrin superfamily proteins differentially bundle actin and localize to spines in CA1 pyramidal neurons. Two-dimensional projections of confocal images of neurons expressing GFP (*A*) or the ABDs of different proteins fused to DsRed (*B–F*). (*B*) Neurons expressing the ABD of β I-spectrin show actin bundling in the cell soma and in dendrites. The ABD of utrophin (*C*) also bundles actin in the soma. Neurons expressing the ABD of actinin (*D*) did not display actin bundling in the soma, dendrites, or spines (Scale right = 25 μ m, Scale left = 100 μ m). *E*, Merged image of a dendrite from a CA1 pyramidal neuron co-expressing actin-GFP and the ABD of β I-spectrin fused to DsRed. The arrow indicates a bundled actin filament in the dendrite. Images at the right show the co-localization of actin and the ABD of β I-spectrin in spine heads (scale bar = 1 μ m). *F*, Co-expression of the ABDs of α -actinin, utrophin, dystrophin, or filamin, all fused to DsRed, and GFP in CA1 pyramidal cells demonstrate their differential targeting to dendritic spines (*E*, *F* Scale bar = 10 μ m). doi:10.1371/journal.pone.0016197.g001

but significant, decrease in average spine density as compared to GFP transfected neurons ($n = 5$ neurons/group $p < 0.01$, ANOVA and Tukey's test)(Fig. 3C).

Only the ABD of β I-spectrin affected dendritic spines. Dendritic spine heads in cells expressing GFP plus the ABDs of α -actinin-2,

dystrophin, utrophin and filamin were not significantly larger than spines in cells expressing GFP alone ($n = 5$ neurons/group $p > 0.05$, ANOVA and Tukey's test)(Fig. 2C). Because all of these constructs bind actin, some other action of the β I-spectrin ABD must be responsible for the enlargement of the dendritic spine head.




Figure 2. The ABD of β -spectrin specifically enlarges dendritic spine heads. *A*, Images of GFP fluorescence in cells co-transfected with GFP and the ABD of β -spectrin fused to DsRed, or with GFP alone, illustrate the substantial increase in spine head diameter seen when the ABD of β -spectrin is overexpressed. Scale = 4 μ m (left) and 2 μ m (right). *B*, Cumulative probability plots demonstrate the significant increase in spine head diameter in cells expressing ABD of β -spectrin when compared to cells transfected with GFP alone, and the lack of a significant effect on spine length ($n=500$ spines/5 neurons/group; $*=p<0.05$, K-S test). *C*, When compared to the ABD of other spectrin superfamily proteins, only the spines expressing the ABD of β -spectrin were significantly enlarged ($n=5$ neurons/construct; $*=p<0.01$, ANOVA with Tukey's test). doi:10.1371/journal.pone.0016197.g002




Figure 3. The ABD of β I-spectrin prevents actin depolymerization. **A**, After application of latrunculin ($5 \mu\text{M}$) for 5 hr, the diameter of the spine head in cells expressing GFP alone decreased significantly ($n = 500$ spines/5 neurons/group; $p < 0.05$, K-S test), whereas spine length was not changed. **B**, Latrunculin treatment produced no significant decrease in spine head diameter or spine length in cells co-expressing the ABD of β I-spectrin fused to DsRed and GFP, in contrast. Scale bar = $2 \mu\text{m}$ for **A** and **B**. **C**, Application of latrunculin-B resulted in a significant decrease in spine density in spines expressing GFP alone ($n = 5$ neurons/group; $p < 0.01$ ANOVA with Tukey's test), but not in spines expressing the ABD of β I-spectrin. Spine density was significantly lower in cells expressing the β I-spectrin ABD, as compared to cells expressing GFP alone. Scale bar = $6 \mu\text{m}$. doi:10.1371/journal.pone.0016197.g003

The ABD of β I-spectrin stabilizes actin filaments in dendritic spines

Differences in the sizes of spine heads correlate with different states of actin polymerization [38]. We further probed the activity of the ABD of β I-spectrin on actin by testing its ability to stabilize actin filaments in spines in the presence of latrunculin-B, which depolymerizes F-actin by preventing addition of G-actin monomers [15,39]. F-actin in spines is significantly depolymerized after 1 hour of latrunculin treatment, with complete depolymerization requiring 4–8 h. This is accompanied by the collapse of the dendritic spine head, known as “deflation” [15,21,39]. In agreement with these

results, we found that application of latrunculin-B ($6 \mu\text{M}$) for 5 h to slice cultures caused a 50% decrease in spine head diameter in cells transfected with GFP alone (GFP+latrunculin = $0.4 \mu\text{m} \pm 0.1 \mu\text{m}$ vs. GFP alone = $0.8 \mu\text{m} \pm 0.1 \mu\text{m}$; $n = 500$ spines/5 neurons/group; $p < 0.05$, K-S test)(Fig. 3A). Spine length was not affected significantly, however ($1.2 \mu\text{m} \pm 0.06 \mu\text{m}$ control vs. $1.2 \mu\text{m} \pm 0.04 \mu\text{m}$ latrunculin; $n = 500$ spines/5 neurons/group; $p > 0.05$, K-S test). In contrast, application of latrunculin to cells transfected with the ABD of β I-spectrin had no significant effect on either spine head diameter (ABD of β I-spectrin+latrunculin = $2.0 \mu\text{m} \pm 0.20 \mu\text{m}$ vs. ABD of β I-spectrin = $2.1 \mu\text{m} \pm 0.2 \mu\text{m}$; $n = 500$ spines/5 neurons/group; $p > 0.05$,

K-S test) or spine length ($1.19 \mu\text{m} \pm 0.04 \mu\text{m}$ control vs. $1.2 \mu\text{m} \pm 0.06 \mu\text{m}$ latrunculin; $n = 500$ spines/5 neurons/group; $p > 0.05$, K-S test)(Fig. 3B). Treatment with latrunculin-B also resulted in a significant decrease in overall spine density in GFP expressing neurons ($n = 5$ neurons/group, $p < 0.01$, ANOVA and Tukey's test), but not in neurons expressing the ABD of β I-spectrin ($n = 5$ neurons/group, $p > 0.05$, ANOVA and Tukey's test)(Fig. 3C). Actin in spines expressing the β I-spectrin ABD was thus resistant to the depolymerizing actions of prolonged latrunculin treatment.

We next tested the effects of acute application of latrunculin-B in cells transfected with actin-GFP alone or with the β I-spectrin ABD (Fig. 4). Depolymerization of actin with latrunculin reduces the intensity of actin-GFP fluorescence in the spine head as the actin monomers that cannot polymerize diffuse out of the spine [40]. We found that acute addition of latrunculin-B ($5 \mu\text{M}$) induced a rapid (< 3 min) decrease in actin-GFP intensity in control cells, as previously reported. Latrunculin application had a significantly smaller effect on the intensity of actin-GFP in spines of cells co-transfected with the ABD of β I-spectrin ($n = 5$ neurons/group, $p < 0.001$, repeated measures ANOVA). This effect was specific for the ABD of β I-spectrin, because latrunculin produced a strong decrease in the fluorescence intensity of spines co-transfected with the ABD of α -actinin-2 and actin-GFP, comparable to its effects on cells expressing actin-GFP alone. We conclude that the presence of the ABD of β I-spectrin in spines stabilizes F-actin and reduces its rate of depolymerization.

The ABD of β I-spectrin decreases the morphological plasticity of dendritic spines

Dynamic morphological changes in the shape of dendritic spine heads ("morphing") occur on the order of minutes or seconds and are dependent upon actin polymerization and depolymerization in the spine head [10,40]. We therefore tested if decreases in actin depolymerization resulting from the expression of the ABD of β I-spectrin would decrease dendritic spine motility. We used live-cell imaging to compare the spontaneous, constitutive morphing of dendritic spines in neurons transfected with GFP alone or GFP together with the ABD of β I-spectrin (Fig. 5). Because the activity of both AMPARs and NMDARs may affect spine morphing [10], we also tested the effects of applying saturating concentrations of DNQX ($40 \mu\text{M}$) and AP5 ($40 \mu\text{M}$). Using GFP emission, we measured the variance of spine head diameter over a period of 10 min to quantify spine head dynamics and to control for differences in spine head size induced by the constructs. Spines expressing GFP alone exhibited significantly more spontaneous morphing than spines expressing the ABD of β I-spectrin ($p < 0.001$, Kruskal-Wallis test), and neither were affected by DNQX and AP5 (Fig. 5C). We conclude that stabilization of actin filaments by the ABD of β I-spectrin attenuates the structural dynamics of dendritic spines.

Functional consequences of expressing the ABD of β I-spectrin

We next asked if there were functional correlates of the morphological changes induced by the ABD of spectrin. The postsynaptic density (PSD) is a specialized structure at the tip of the dendritic spine head comprised of densely packed glutamate receptors and signaling proteins [41,42] and its size is positively correlated with synaptic strength [2]. A principal constituent of the PSD is the scaffolding protein, PSD-95 [43]. As a first test of the possibility that the enlargement of spine heads induced by the ABD of β I-spectrin (lacking the DsRed tag) would be associated with an increase in synaptic strength, we co-expressed it with GFP-

tagged PSD-95 and cytosolic mCherry (Fig. 6A). We observed a significant increase in the area of PSD-95-GFP in dendritic spines co-expressing the ABD of β I-spectrin, as compared to cells expressing PSD-95-GFP alone (mean area = $0.20 \mu\text{m}^2 \pm 0.02 \mu\text{m}^2$ in controls vs. $0.38 \mu\text{m}^2 \pm 0.01 \mu\text{m}^2$ with the ABD of β I-spectrin; $n = 365$ spines/4 neurons/group; $p < 0.05$, K-S test)(Fig. 6B). In agreement with previous reports [44] we observed a linear relationship between spine head size and PSD-95 area under both conditions ($r = 0.83$ for control and 0.79 for the ABD of β I-spectrin). We conclude that overexpression of the ABD of β I-spectrin is associated with an increase in the size of the PSD.

It has been demonstrated that the number of AMPARs is proportional to PSD area [2,45,46]. We therefore asked whether the increase in PSD size produced by the ABD of β I-spectrin would be accompanied by an increase in the number of synaptic AMPARs in the spine head. To answer this question, we transfected CA1 pyramidal cells in hippocampal slice cultures with the ABD of β I-spectrin tagged with DsRed and performed whole-cell voltage-clamp recordings of spontaneous miniature excitatory postsynaptic currents (mEPSCs) on transfected and untransfected neurons in the presence of TTX ($1 \mu\text{M}$). Alexa 488 was included in the pipette solution in order to ensure that the recorded neurons were truly those expressing the spectrin construct, as apparent by green and red fluorescence. Mean mEPSC amplitudes were significantly larger in cells transfected with the ABD of β I-spectrin compared to untransfected cells in the same cultures, consistent with an increase in the number of synaptic AMPARs (mean amplitude = 20.8 ± 1.4 pA for cells expressing the ABD of β I-spectrin vs. 17.1 ± 1.1 pA in untransfected cells; $n = 7$ neurons/group; $p < 0.05$, K-S test)(Fig. 7A,B). The frequency of mEPSCs was decreased $\sim 20\%$ after transfection with the ABD of β I-spectrin, compared to untransfected neurons recorded from the same slice ($p < 0.05$, t-test)(Fig. 7C), consistent with the decrease in spine density described above. To test for non-specific effects of biolistic transfection on neuronal function, we also compared mEPSC amplitudes in neurons transfected with GFP alone and untransfected cells and observed no significant difference (mean amplitude = 17.1 ± 0.5 pA in untransfected cells vs. 15.7 ± 0.4 pA in cells expressing GFP; $n = 7$ neurons/group, $p > 0.05$, unpaired t-test). The presence of the ABD of β I-spectrin in dendritic spines thus increased the sizes of the PSD and the spine head, as well as synaptic strength.

The activity of the ABD of β I-spectrin is mediated by Rac3

Do the effects of the β I-spectrin ABD on spines result directly from its interaction with actin or do they result because of recruitment of some other molecule to the spine head? Our observation that only the ABD of β I-spectrin localizes to spines and caused their enlargement, although all of them bind to actin (Fig. 2), favors the latter hypothesis. Small GTPases are important signaling molecules involved in the regulation of dendritic spine morphological dynamics [47]. One small GTPase, Rac1, modulates dendritic spine morphology [48] and function [49]. We therefore tested the role of rac proteins in the morphological changes induced by the ABD of β I-spectrin.

Two Rac GTPases are expressed in hippocampal and neocortical pyramidal cells, Rac1 and Rac3 [50,51]. We first immunoprecipitated endogenous β I-spectrin from cultured cortical neurons and screened for the presence of Rac3 and Rac1. Surprisingly, Rac3 was enriched in immunoprecipitates of β I-spectrin, but Rac1 was absent (3 replicates)(Fig. 8A). We tested the importance of this interaction by co-expressing a dominant negative Rac3 construct together with the ABD of β I-spectrin. Under these conditions, spine head diameters were significantly




Figure 4. The ABD of β -spectrin prevents acute actin depolymerization. *A*, Images of GFP fluorescence in spines from CA1 pyramidal neurons expressing actin-GFP alone (upper row), or co-expressing actin-GFP together with the ABD of β -spectrin (lower row) before and after the application of 5 μ M latrunculin-B. Note that fluorescence decreased within 5 minutes of latrunculin treatment in cells expressing GFP, but not in cells co-expressing the spectrin ABD. Scale bar = 2 μ m. *B*, Normalized fluorescence emission of dendritic spines was imaged every 30 seconds before and after the addition of latrunculin-B. Fluorescence decreased rapidly (<3 min) in spines expressing actin-GFP alone and in spines in which the ABD of α -actinin was co-expressed, but not in spines co-expressing the ABD of β -spectrin and actin-GFP ($n = 5$ neurons/construct; $p < 0.001$, repeated measures ANOVA).
doi:10.1371/journal.pone.0016197.g004

smaller than in cells expressing the β I-spectrin ABD alone (Fig. 8B,C). Spine head diameters in cells co-expressing the β I-spectrin ABD and dominant negative Rac3 or Rac3-DN alone

were not significantly different than in cells expressing only GFP, suggesting that Rac3 signaling is required for the effects of the β I-spectrin ABD on spines. If so, then this effect should be mimicked




Figure 5. The ABD of β -spectrin decreases the morphological plasticity of dendritic spines. *A*, Time-lapse images in control saline of GFP fluorescence in a single spine from a CA1 pyramidal neuron expressing either GFP alone (upper row) or GFP together with the ABD of β -spectrin (lower row). Rapid, ongoing changes in the shape of the spine head occurred in spines expressing GFP alone, but not in spines expressing both GFP and the ABD of β -spectrin. Scale bar = 1 μm ; time in sec. *B*, Changes in spine head diameter in the spines from *A* were plotted over time. Spines expressing GFP exhibited more spontaneous morphing, compared to a spine expressing the ABD of β -spectrin, regardless of whether NMDARs and AMPARs were active. *C*, Group data showing the variance of spine head diameter in neurons expressing GFP alone or the ABD of β -spectrin in the

two conditions. Spines expressing the ABD of β I-spectrin displayed significantly less variance in the shape of their heads than did spines expressing GFP alone ($n=5$ neurons/group; $*=p<0.001$, Kruskal-Wallis test). There was no significant effect of blocking NMDARs and AMPARs on spontaneous morphing in any condition.
doi:10.1371/journal.pone.0016197.g005

by over-expressing wild-type Rac3. Indeed, the size of dendritic spine heads in CA1 pyramidal cells transfected with Rac3 was comparable to spines in cells expressing the ABD of β I-spectrin. Overexpression of Rac3 also resulted in an increase in mEPSC amplitude. The mean amplitude of mEPSCs recorded from neurons expressing Rac3 was not significantly different from those expressing the ABD of β I-spectrin and both were larger than for mEPSCs in untransfected or GFP-transfected cells (Fig. 7B). We conclude that Rac3 can alter dendritic spine structure and function in hippocampal pyramidal cells.

Discussion

Actin is highly concentrated in dendritic spines and determines their morphology [52–54]. The actin cytoskeleton provides a scaffold to which a variety of transmembrane proteins and their intracellular signaling partners are anchored, often indirectly via actin binding proteins. The spectrin superfamily of proteins can form complexes via their actin binding domains, which interact with a number of other proteins, such as RACK-1, protein 4.1 and dynactin [25,55,56]. β I-spectrin has been immunocytochemically




Figure 6. Expression of the ABD of β I-spectrin increases the size of the PSD. **A**, Merged images from CA1 pyramidal neurons in hippocampal slice cultures co-transfected with PSD-95 fused to GFP (green) and mCherry (red), with or without an unlabelled β I-spectrin ABD construct. The areas of the clusters of PSD-95 in the heads of spines expressing the ABD of β I-spectrin were larger than in cells expressing PSD-95 alone. Scale bar = 4 μ m (left), 1 μ m (right). **B**, Cumulative probability plot showing that the area of the PSD was significantly larger in spines expressing PSD-95-GFP together with the ABD of β I-spectrin than in spines expressing PSD-95-GFP alone ($n=400$ spines/4 neurons/condition; $p<0.05$, K-S test).
doi:10.1371/journal.pone.0016197.g006




Figure 7. The ABD of β -spectrin enhances single spine AMPAR-mediated synaptic responses. *A*, mEPSCs from an untransfected control cell (upper trace) and a cell transfected with the ABD of β -spectrin (lower trace) were recorded at -70 mV in the presence of TTX ($1 \mu\text{M}$). Average sized single mEPSCs recorded from the untransfected cell (black) and transfected cell (gray) are shown superimposed at right. *B*, Group data showing that neurons expressing the ABD of β -spectrin had significantly greater mEPSC amplitudes, compared to untransfected controls or cells expressing GFP alone, suggesting an increase in AMPARs at these synapses ($n = 7$ neurons/group; $p < 0.05$, K-S test). The increase in mEPSC amplitude produced by the spectrin ABD was mimicked in cells expressing CA Rac3, in which mEPSC amplitude was also larger than in controls or GFP expressing cells ($p < 0.05$, K-S test). *C*, The frequency of mEPSCs in transfected neurons decreased significantly, suggesting fewer functional synapses ($n = 7$ neurons/group; $* = p < 0.05$, t-test). doi:10.1371/journal.pone.0016197.g007

detected in dendritic spines, but its functions there remain unknown.

In this study, we expressed the actin binding domain of β -spectrin and related spectrin superfamily proteins, an approach used successfully as a first step in revealing the function of the full-length proteins, such as α -actinin [30]. Our assumption is that the exogenous ABD construct will mimic the function of the N-terminal ABD of the full length protein. We cannot exclude, however, that overexpression of the ABD of β -spectrin may negatively affect endogenous spectrin or induce other downstream compensatory effects [30]. Indeed, a full characterization of spectrin requires complimentary approaches, such as knock-down of endogenous spectrin and overexpression of the full length protein, that are beyond the scope of the current study. Nevertheless, spectrin is a prominent substrate of the Ca^{2+} -activated protease calpain and activation of calpain may result in the creation of fragments not unlike those we have expressed.

Interactions between β -spectrin and actin

Although all of the proteins of the spectrin superfamily and related proteins share highly homologous ABDs, they are all unique at many sites (Fig. S1). Indeed, we found a surprising heterogeneity in the way these ABDs were targeted in CA1 pyramidal cells. The ABDs of β -spectrin and utrophin, but not dystrophin, colocalized with GFP-tagged actin and bundled actin filaments in the cell body and proximal dendrites. The ABDs of β -spectrin, α -actinin-2 and filamin, but not utrophin and dystrophin, were concentrated with actin in the heads of dendritic spines. Our results with the ABD of α -actinin-2 are in contrast to those of Nakagawa et al. [30], who reported that the ABD of α -actinin-2 did not localize to spines, although comparable regions of the protein were used in both studies. Even more unique to the ABD of β -spectrin was its ability to enlarge the spine head. None of the other ABDs we assayed produced this effect. Nakagawa et al. [30] observed that overexpression of full length α -actinin-2




Figure 8. Rac3 is immunoprecipitated with the ABD of β -spectrin and might mediate its effects on dendritic spines. *A*, Rac3 is specifically enriched in immunoprecipitates of detergent extracts of cortical cell cultures generated with antibodies to β -spectrin. The input sample (lane 1) contained easily detected amounts of β -spectrin and Rac1 but relatively little Rac3. Immunoprecipitation with an anti- β -spectrin antibody (lane 2) specifically concentrated β -spectrin and Rac3, but did not precipitate Rac1. Neither GTPase was detected after immunoprecipitation with a non-immune rabbit IgG (lane 3). *B*, GFP images of CA1 pyramidal cell dendrites co-expressing GFP and the ABD of β -spectrin either with or without dominant negative (DN) Rac3, and co-expressing GFP with wild type Rac3. Scale bar = 10 μm . *C*, Pooled data illustrate the effects on the mean spine head diameter, calculated from GFP images ($n=100\text{--}300$ spines/neuron, 2–5 neurons/group, $p<0.001$, ANOVA with Sheffe's test). Spines in cells expressing the ABD of β -spectrin had larger heads than cells expressing GFP alone, but not when co-expressing dominant negative Rac3. Expression of wild-type Rac3 resulted in spines with large heads, thus mimicking the effects of the ABD of β -spectrin. doi:10.1371/journal.pone.0016197.g008

elongated dendritic spines and that this required its ABD. These results suggest that different actin-binding proteins from the spectrin superfamily can exert differential structural effects on dendritic spines, but only the ABD of β -spectrin can enlarge postsynaptic spine heads.

The ABD of β -spectrin also affected the stability of actin filaments within the spine. In control cells, both acute (minutes) and chronic (5 hr) application of latrunculin-B, an inhibitor of actin polymerization, reduced the dimensions of spine heads and caused a loss of GFP-actin from spines, as reported previously [40]. In cells expressing the ABD of β -spectrin, in contrast, both

the latrunculin-induced loss of actin-GFP and deflation of the spine head were inhibited. Thus, the ABD of β -spectrin reduces the rate at which actin filaments are depolymerized, even when the addition of new actin monomers has been blocked by latrunculin-B. This effect was also specific, as the ABD of α -actinin-2 did not stabilize actin-GFP in spines. These data suggest that some of the effects of expressing these constructs result not from their ability to bind to actin, but perhaps because they recruit other signaling molecules differentially.

Consistent with its ability to stabilize actin filaments, the ABD of β -spectrin significantly inhibited the constitutive morphing of

dendritic spines. Prevention of actin polymerization with latrunculin-B or cytochalasin-D has been reported previously to eliminate dynamic changes in spine head shape [10]. We suggest that endogenous β I-spectrin in spines maintains the spine head in a large and stable state, in part by stabilizing filamentous actin, and that decreases in β I-spectrin might facilitate actin reorganization and promote dynamic changes in spine size and shape.

Functional consequences of expressing the ABD of β I-spectrin

The turnover of F-actin in dendritic spines is necessary for postsynaptic reorganization and the translocation of proteins into and out of the synaptic plasma membrane [14,58]. This activity is driven by rapid actin polymerization and depolymerization [59]. For example, inhibition of actin polymerization by latrunculin-A or ADF/cofilin increases the relative rate of AMPAR endocytosis and partially inhibits late-phase LTP [16,22]. Conversely, the activity of glutamate receptors can affect actin turnover [60] by modulating actin polymerization or depolymerization. Indeed, another spectrin superfamily protein, α -actinin-2, regulates not only the morphology of dendritic spines [57], but also the trafficking of AMPARs [61] and the Ca^{2+} -dependent inactivation of NMDARs [62,63].

Larger dendritic spines contain more AMPARs and generate larger responses to glutamate [2,46]. We observed an increase in the size of dendritic spine heads after expressing the ABD of β I-spectrin, as well as an increase in the area occupied by PSD-95 in the spine head. Stabilization of F-actin by the β I-spectrin ABD, which increases the size of spine heads, is likely to account for the increase in PSD size [58]. The association of PSD-95 with F-actin is required to anchor AMPARs at postsynaptic sites [64]. We observed that the amplitudes of AMPAR-mediated mEPSCs were significantly larger in cells expressing the ABD of β I-spectrin. The doubling in PSD-95 area and spine head diameter observed in cells expressing the β I-spectrin ABD was associated with only a $\sim 40\%$ increase in mean mEPSC amplitude, however, indicating that a strict correlation between postsynaptic density size and AMPA receptor number [e.g. 46] is not obligatory under all conditions. Nevertheless, our data represent the first evidence that the actin-binding activity of proteins of the spectrin superfamily can affect postsynaptic sensitivity to released neurotransmitter.

Rac3 and the ABD of β I-spectrin

How might endogenous β I-spectrin regulate synaptic function? In addition to its ABD, β I-spectrin contains binding sites for calmodulin, protein 4.1, ankyrin, and other proteins [25,65,66]. Protein 4.1 promotes the association of the ABD of spectrin with F-actin [67,68]. The neuronal splice form, 4.1N, interacts with AMPA-type GluR1 subunits and promotes their surface expression [69]. Recruitment of protein 4.1 by the β I-spectrin ABD may therefore promote the insertion of AMPARs into the postsynaptic membrane.

Small GTPases such as Rho, Rac and cdc42, promote morphological plasticity by regulating dynamic changes in the actin cytoskeleton. For example, constitutively active Rac1 disorganizes spines [70], whereas dominant-negative Rac1 disrupts spines [47,70]. Rac1 activity increases following activation of synaptic NMDARs and promotes the subsequent enlargement of dendritic spines [49]. Karilin-7, a guanine nucleotide exchange factor specific for Rac1, is targeted to dendritic spines, where it enlarges spine heads [71] and potentiates AMPAR-mediated synaptic activity [49]. Less is known about Rac3, although it is highly homologous to Rac1 and is expressed in neurons [51], particularly in the CA1 and CA3 region of the hippocampus during development [50]. Rac3 promotes neurite extension in cultured hippocampal neurons [72] and induces effects on cell

differentiation and morphology in neuronal cell lines that are distinct from those caused by Rac1 [73]. Mice in which Rac3 is genetically deleted are viable, learn a hippocampal dependent task normally, and have normal dendritic spine densities, although this may be due to compensation by Rac1 [74,75].

We observed that Rac3 co-immunoprecipitates with spectrin, but not Rac1. Overexpression of wild-type Rac3 increased the sizes of spine heads as effectively as overexpression of the ABD of β I-spectrin. In contrast, co-expression of a dominant-negative form of Rac3 with the ABD of β I-spectrin with inhibited its ability to enhance spine head size. Finally, neurons that overexpress Rac3 had increased mEPSC amplitudes. We therefore suggest that Rac3 activation is promoted in part by the ABD of β I-spectrin and may mediate some of the effects of endogenous spectrin on postsynaptic structure and function.

How might Rac3 mediate its effects? Rac3 associates with neurabin and this association is required for its GTPase activity [72]. Furthermore, neurabin binds to actin [76] and this binding is required for its localization in dendritic spines [39]. Overexpression of neurabin increases spine head size [77] and AMPAR expression and triggers actin bundling [78]. It is thus possible that the presence of high levels of β I-spectrin ABD in the spine head leads to spine enlargement and synapse strengthening by recruiting Rac3 and neurabin. Alternatively, neurabin may exert its effects on dendritic spines [39] by binding to actin and, like β I-spectrin, recruiting Rac3. It should be noted that the dominant negative Rac3 construct did not *reduce* dendritic spine size compared to control cells. Similarly, genetic deletion of Rac3 does not affect dendritic spine density in hippocampal cell cultures [75], although spine size was not quantified. This suggests that the interaction between endogenous β I-spectrin and Rac3 is not necessary to *maintain* spine head size. This possibility could be tested in future studies in which expression of endogenous β I-spectrin is reduced by targeted siRNA.

Our results suggest the testable hypothesis that the ABD of endogenous β I-spectrin and Rac3 can interact in a complex that promotes both activation of Rac3 and polymerization of actin. Changes in the expression of β I-spectrin might thus affect the dynamic state of the actin cytoskeleton in spines, thereby influencing AMPAR trafficking and the balance between synapse plasticity and stability.

Supporting Information

Figure S1 Sequence alignment of spectrin family ABDs.

Although the sequences are identical to β I-spectrin at many residues (indicated by *), and very similar (indicated by :) or homologous at others (indicated by.), there are many differences. Although all share the ability to bind and bundle actin, these regions of difference are presumably responsible for the distinct activities observed in the various experiments. (TIF)

Acknowledgments

We thank Ms. Wendy Resneck for her expert assistance with immunoprecipitations, Drs. T. Abrams and T. Blanpied for comments on the manuscript, and Dr. T. Blanpied and Ms. E. Kong for providing the cortical cell cultures. Portions of these data have been presented to the Society for Neuroscience in 2005.

Author Contributions

Conceived and designed the experiments: MWN RJB SMT. Performed the experiments: MWN XC. Analyzed the data: MWN. Contributed reagents/materials/analysis tools: MRS. Wrote the paper: MWN RJB SMT.

References

- Ziv NE, Smith SJ (1996) Evidence for a role of dendritic filopodia in synaptogenesis and spine formation. *Neuron* 17: 91–102.
- Matsuzaki M, Ellis-Davies GC, Nemoto T, Miyashita Y, Iino M, et al. (2001) Dendritic spine geometry is critical for AMPA receptor expression in hippocampal CA1 pyramidal neurons. *Nat Neurosci* 4: 1086–1092.
- Kopec CD, Li B, Wei W, Boehm J, Malinow R (2006) Glutamate receptor exocytosis and spine enlargement during chemically induced long-term potentiation. *J Neurosci* 26: 2000–2009.
- Zito K, Scheuss V, Knott G, Hill T, Svoboda K (2009) Rapid functional maturation of nascent dendritic spines. *Neuron* 61: 247–258.
- Kozorovitskiy Y, Gross CG, Kopil C, Battaglia L, McBreen M, et al. (2005) Experience induces structural and biochemical changes in the adult primate brain. *Proc Natl Acad Sci USA* 102: 17478–17482.
- Trachtenberg JT, Chen BE, Knott GW, Feng G, Sanes JR, et al. (2002) Long-term in vivo imaging of experience-dependent synaptic plasticity in adult cortex. *Nature* 420: 788–794.
- Matsuzaki M, Honkura N, Ellis-Davies GC, Kasai H (2004) Structural basis of long-term potentiation in single dendritic spines. *Nature* 429: 761–766.
- Nägerl UV, Eberhorn N, Cambridge SB, Bonhoeffer T (2004) Bidirectional activity-dependent morphological plasticity in hippocampal neurons. *Neuron* 44: 759–767.
- Bagal AA, Kao JP, Tang CM, Thompson SM (2005) Long-term potentiation of exogenous glutamate responses at single dendritic spines. *Proc Natl Acad Sci USA* 102: 14434–14439.
- Fischer M, Kaech S, Knutti D, Matus A (1998) Rapid actin-based plasticity in dendritic spines. *Neuron* 20: 847–854.
- Halpain S, Hipolito A, Saffer L (1993) Regulation of F-actin stability in dendritic spines by glutamate receptors and calcineurin. *J Neurosci* 13: 9835–9844.
- Blanpied TA, Kerr JM, Ehlers MD (2008) Structural plasticity with preserved topology in the postsynaptic protein network. *Proc Natl Acad Sci USA* 34: 12587–12592.
- Kim CH, Lisman JE (1999) A role of actin filament in synaptic transmission and long-term potentiation. *J Neurosci* 19: 4314–4324.
- Krucker T, Siggins GR, Halpain S (2000) Dynamic actin filaments are required for stable long-term potentiation (LTP) in area CA1 of the hippocampus. *Proc Natl Acad Sci USA* 97: 6856–6861.
- Zhang W, Benson DL (2000) Development and molecular organization of dendritic spines and their synapses. *Hippocampus* 10: 512–526.
- Fukazawa Y, Saitoh Y, Ozawa F, Ohta Y, Mizuno K, et al. (2003) Hippocampal LTP is accompanied by enhanced F-actin content within the dendritic spine that is essential for late LTP maintenance in vivo. *Neuron* 38: 447–460.
- Okamoto K, Nagai T, Miyawaki A, Hayashi Y (2004) Rapid and persistent modulation of actin dynamics regulates postsynaptic reorganization underlying bidirectional plasticity. *Nat Neurosci* 7: 1104–1112.
- Hering H, Sheng M (2003) Activity-dependent redistribution and essential role of cortactin in dendritic spine morphogenesis. *J Neurosci* 23: 11759–11769.
- Ethell IM, Pasquale EB (2005) Molecular mechanisms of dendritic spine development and remodeling. *Prog Neurobiol* 75: 161–205.
- Bramham CR (2008) Local protein synthesis, actin dynamics, and LTP consolidation. *Curr Opin Neurobiol* 18: 524–531.
- Allison DW, Gelfand VI, Spector I, Craig AM (1998) Role of actin in anchoring postsynaptic receptors in cultured hippocampal neurons: differential attachment of NMDA versus AMPA receptors. *J Neurosci* 18: 2423–2436.
- Zhou Q, Xiao M, Nicoll RA (2001) Contribution of cytoskeleton to the internalization of AMPA receptors. *Proc Natl Acad Sci USA* 98: 1261–1266.
- Davis LH, Bennett V (1994) Identification of two regions of beta G spectrin that bind to distinct sites in brain membranes. *J Biol Chem* 269: 4409–4416.
- Bennett V, Healy J (2008) Organizing the fluid membrane bilayer: diseases linked to spectrin and ankyrin. *Trends Mol Med* 14: 28–36.
- Bennett V, Baines AJ (2001) Spectrin and ankyrin-based pathways: metazoan inventions for integrating cells into tissues. *Physiol Rev* 81: 1353–1392.
- Zhou M, Baudry M (2006) Developmental changes in NMDA neurotoxicity reflect developmental changes in subunit composition of NMDA receptors. *J Neurosci* 26: 2956–2963.
- Calvert R, Bennett P, Gratzner W (1980) Properties and structural role of the subunits of human spectrin. *Eur J Biochem* 107: 355–361.
- Li X, Bennett V (1996) Identification of the spectrin subunit and domains required for formation of spectrin/adducin/actin complexes. *J Biol Chem* 271: 15695–15702.
- Ursitti JA, Martin L, Resneck WG, Chaney T, Zielke C, et al. (2001) Spectrins in developing rat hippocampal cells. *Brain Res Dev Brain Res* 129: 81–93.
- Nakagawa T, Engler JA, Sheng M (2004) The dynamic turnover and functional roles of alpha-actinin in dendritic spines. *Neuropharmacology* 47: 734–745.
- Gähwiler BH, Thompson SM, McKinney RA, Debanne D, Robertson RT (1998) Organotypic slice cultures of neural tissue. In: *Culturing Nerve Cells* (2nd Ed.), G Banker, K Goslin, eds. MIT Press: Cambridge, MA, pp 461–498.
- Stone MR, O'Neill A, Catino D, Bloch RJ (2005) Specific interaction of the actin-binding domain of dystrophin with intermediate filaments containing keratin 19. *Mol Biol Cell* 16: 4280–4293.
- Amann KJ, Renley BA, Ervasti JM (1998) A cluster of basic repeats in the dystrophin rod domain binds F-actin through an electrostatic interaction. *J Biol Chem* 273: 28419–28423.
- Galkin VE, Orlova MS, VanLoock IN (2002) The utrophin actin-binding domain binds F-actin in two different modes: implications for the spectrin superfamily of proteins. *J Cell Biol* 157: 243–251.
- McAllister AK (2000) Biolistic Transfection of Neurons. *Sci STKE* 2000, p11.
- Oray S, Majewska A, Sur M (2006) Effects of synaptic activity on dendritic spine motility of developing cortical layer v pyramidal neurons. *Cereb Cortex* 16: 730–741.
- Zhou D, Ursitti JA, Bloch RJ (1998) Developmental expression of spectrins in rat skeletal muscle. *Mol Biol Cell* 9: 47–61.
- Capani F, Martone ME, Deerinck TJ, Ellisman MH (2001) Selective localization of high concentrations of F-actin in subpopulations of dendritic spines in rat central nervous system: a three-dimensional electron microscopic study. *J Comp Neurol* 435: 156–170.
- Zito K, Knott G, Shepherd GM, Shenolikar S, Svoboda K (2004) Induction of spine growth and synapse formation by regulation of the spine actin cytoskeleton. *Neuron* 44: 321–334.
- Star EN, Kwiatkowski DJ, Murthy VN (2002) Rapid turnover of actin in dendritic spines and its regulation by activity. *Nat Neurosci* 5: 239–246.
- Okabe S (2007) Molecular anatomy of the postsynaptic density. *Mol Cell Neurosci* 34: 503–518.
- Feng W, Zhang M (2009) Organization and dynamics of PDZ-domain-related supramodules in the postsynaptic density. *Nat Rev Neurosci* 10: 87–99.
- Cho KO, Hunt CA, Kennedy MB (1992) The rat brain postsynaptic density fraction contains a homolog of the Drosophila discs-large tumor suppressor protein. *Neuron* 9: 929–942.
- Harris KM, Stevens JK (1989) Dendritic spines of CA1 pyramidal cells in the rat hippocampus: serial electron microscopy with reference to their biophysical characteristics. *J Neurosci* 9: 2982–2997.
- Takumi Y, Ramirez-Leon V, Laake P, Rinivik E, Ottersen OP (1999) Different modes of expression of AMPA and NMDA receptors in hippocampal synapses. *Nat Neurosci* 2: 618–624.
- Nusser Z, Lujan R, Laube G, Roberts JD, Molnar E, et al. (1998) Cell type and pathway dependence of synaptic AMPA receptor number and variability in the hippocampus. *Neuron* 21: 545–559.
- Nakayama AY, Harms MB, Luo L (2000) Small GTPases Rac and Rho in the maintenance of dendritic spines and branches in hippocampal pyramidal neurons. *J Neurosci* 20: 5329–5338.
- Sin WC, Haas K, Ruthazer ES, Cline HT (2002) Dendrite growth increased by visual activity requires NMDA receptor and Rho GTPases. *Nature* 419: 475–480.
- Xie Z, Srivastava DP, Photowala H, Kai L, Cahill ME, et al. (2007) Kalirin-7 controls activity-dependent structural and functional plasticity of dendritic spines. *Neuron* 56: 640–656.
- Corbetta S, Gualdoni S, Albertinazzi C, Paris S, Croci L, et al. (2005) Generation and characterization of Rac3 knockout mice. *Mol Cell Biol* 25: 5763–5776.
- Malosio ML, Gilardelli D, Paris S, Albertinazzi C, de Curtis I (1997) Differential expression of distinct members of Rho family GTP-binding proteins during neuronal development: identification of Rac1B, a new neural-specific member of the family. *J Neurosci* 17: 6717–6728.
- Fifkova E, Delay RJ (1982) Cytoplasmic actin in neuronal processes as a possible mediator of synaptic plasticity. *J Cell Biol* 95: 345–350.
- Matus A, Ackermann M, Pehling G, Byers HR, Fujiwara K (1982) High actin concentrations in brain dendritic spines and postsynaptic densities. *Proc Natl Acad Sci USA* 79: 7590–7594.
- Markham JA, Fifkova E (1986) Actin filament organization within dendrites and dendritic spines during development. *Brain Res* 392: 263–269.
- Bennett V, Healy J (2009) Membrane domains based on ankyrin and spectrin associated with cell-cell interactions. *Cold Spring Harb Perspect Biol* 1: 1–17.
- Rodriguez MM, Ron D, Touhara K, Chen CH, Mochly-Rosen D (1999) RACK-1, a protein kinase C anchoring protein, coordinates the binding of activated protein kinase C and select pleckstrin homology domains in vitro. *Biochemistry* 38: 13787–13794.
- Matsuoka Y, Li X, Bennett V (1998) Adducin is an in vivo substrate for protein kinase C: phosphorylation in the MARCKS-related domain inhibits activity in promoting spectrin-actin complexes and occurs in many cells, including dendritic spines of neurons. *J Cell Biol* 142: 485–497.
- Ouyang Y, Wong M, Capani F, Rensing N, Lee CS, et al. (2005) Transient decrease in F-actin may be necessary for translocation of proteins into dendritic spines. *Eur J Neurosci* 22: 2995–3005.
- Pollard TD, Blanchoin L, Mullins RD (2000) Molecular mechanisms controlling actin filament dynamics in nonmuscle cells. *Annu Rev Biophys Biomol Struct* 29: 545–576.
- Fischer M, Kaech S, Wagner U, Brinkhaus H, Matus A (2000) Glutamate receptors regulate actin-based plasticity in dendritic spines. *Nat Neurosci* 3: 887–894.
- Schulz TW, Nakagawa T, Licznernski P, Pawlak V, Kollerker A, et al. (2004) Actin/alpha-actinin-dependent transport of AMPA receptors in dendritic spines: role of the PDZ-LIM protein RIL. *J Neurosci* 24: 8584–8594.
- Rosenmund C, Feltz A, Westbrook GL (1995) Calcium-dependent inactivation of synaptic NMDA receptors in hippocampal neurons. *J Neurophysiol* 73: 427–430.

63. Krupp JJ, Vissel B, Thomas CG, Heinemann SF, Westbrook GL (1999) Interactions of calmodulin and alpha-actinin with the NR1 subunit modulate Ca²⁺-dependent inactivation of NMDA receptors. *J Neurosci* 19: 1165–1178.
64. Lisman JE, Zhabotinsky AM (2001) A model of synaptic memory: a CaMKII/PP1 switch that potentiates transmission by organizing an AMPA receptor anchoring assembly. *Neuron* 31: 191–201.
65. Bennett V, Gilligan DM (1993) The spectrin-based membrane skeleton and micron-scale organization of the plasma membrane. *Annu Rev Cell Biol* 9: 27–66.
66. Viel A, Branton D (1996) Spectrin: on the path from structure to function. *Curr Opin Cell Biol* 8: 49–55.
67. Bennett V, Hughes CA (1995) Adducin: a physical model with implications for function in assembly of spectrin-actin complexes. *Journal of Biol Chem* 270: 18990–18996.
68. Coleman TR, Harris AS, Miche SM, Mooseker MS, Morrow JS (1987) Beta spectrin bestows protein 4.1 sensitivity on spectrin-actin interactions. *J Cell Biol* 104: 519–526.
69. Shen L, Liang F, Walensky LD, Huganir RL (2000) Regulation of AMPA receptor GluR1 subunit surface expression by a 4.1N-linked actin cytoskeletal association. *J Neurosci* 20: 7932–7940.
70. Tashiro A, Minden A, Yuste R (2000) Regulation of dendritic spine morphology by the rho family of small GTPases: antagonistic roles of Rac and Rho. *Cereb Cortex* 10: 927–938.
71. Penzes P, Johnson RC, Sattler R, Zhang X, Huganir RL, et al. (2001) The neuronal Rho-GEF Kalirin-7 interacts with PDZ domain-containing proteins and regulates dendritic morphogenesis. *Neuron* 29: 229–242.
72. Orioli D, Colaluca IN, Stefanini M, Riva S, Dotti CG, et al. (2006) Rac3-induced neuritogenesis requires binding to Neurabin I. *Mol Biol Cell* 17: 2391–2400.
73. Hajdo-Milasinovic, Ellenbroek SIJ, van Es S, van der Vaart B, Collard JG (2007) Rac1 and Rac3 have opposing functions in cell adhesion and differentiation of neuronal cells. *J Cell Sci* 120: 555–566.
74. Corbetta S, D'Adamo P, Gualdoni S, Braschi C, Berardi N, et al. (2008) Hyperactivity and novelty-induced hyperreactivity in mice lacking Rac3. *Behav Brain Res* 186: 246–255.
75. Corbetta S, Gualdoni S, Ciceri G, Monari M, Zuccaro E, et al. (2009) Essential role of Rac1 and Rac3 GTPases in neuronal development. *FASEB J* 23: 1347–1357.
76. Nakanishi H, Obaishi H, Satoh A, Wada M, Mandai K, et al. (1997) Neurabin: a novel tissue-specific actin filament-binding protein involved in neurite formation. *J Cell Biol* 139: 951–961.
77. Oliver CJ, Terry-Lorenzo RT, Elliott E, Bloomer WA, Li S, et al. (2002) Targeting protein phosphatase 1 (PP1) to the actin cytoskeleton: the neurabin I/PP1 complex regulates cell morphology. *Mol Cell Biol* 22: 4690–4701.
78. Terry-Lorenzo RT, Roadcap DW, Otsuka T, Blanpied TA, Zamorano PL, et al. (2005) Neurabin/protein phosphatase-1 complex regulates dendritic spine morphogenesis and maturation. *Mol Biol Cell* 16: 2349–2362.